

Lecture 4: Kernel methods, representer theorem and RKHSs

August 24, 2021

Lecturer: Lei Wu

Scribe: Lei Wu

1 Feature-based methods

We start by considering the linear regression, for which the hypothesis class is

$$\mathcal{F} = \{\beta^T x : \beta \in \mathbb{R}^d\},$$

where we omit the bias term for simplicity. The **ridge regression** penalizes the squared ℓ_2 norm of β :

$$\hat{\beta}_n = \operatorname{argmin}_{\beta \in \mathbb{R}^d} \frac{1}{n} \sum_{i=1}^n (\beta^T x_i - y_i)^2 + \lambda \|\beta\|_2^2.$$

The minimizer has a closed-form solution:

$$\hat{\beta}_n = \left(\frac{1}{n} X^T X + I \right)^{-1} \frac{1}{n} X y,$$

where $X = (x_1, \dots, x_n)^T \in \mathbb{R}^{n \times d}$, $y = (y_1, \dots, y_n)^T \in \mathbb{R}^n$. Another population one is LASSO, which penalizes the ℓ_1 norm of parameters:

$$\min_{\beta \in \mathbb{R}^d} \frac{1}{n} \sum_{i=1}^n (\beta^T x_i - y_i)^2 + \lambda \|\beta\|_1.$$

To consider nonlinear functions, we can consider the model:

$$f(x; \beta) = \sum_{j=1}^m \beta_j \varphi_j(x).$$

Here, $\varphi_1, \dots, \varphi_m$ are a set of (nonlinear) basis functions, which are often referred to as *features* in machine learning. Accordingly, the feature map is defined as $\Phi : \mathcal{X} \mapsto \mathbb{R}^m$ with $\Phi(x) = (\varphi_1(x), \dots, \varphi_m(x))^T \in \mathbb{R}^m$. Typical examples includes

- Spectral methods: $\{\varphi_j\}$ are either Fourier basis or orthogonal polynomials.
- Splines: $\{\varphi_j\}$ are piecewise polynomials.
- Computer vision: Some hand-crafted features.

We can consider two types of feature-based methods.

$$\frac{1}{n} \sum_{i=1}^n (\beta^T \Phi(x_i) - y_i)^2 + \lambda \|\beta\|_2^2$$

$$\frac{1}{n} \sum_{i=1}^n (\beta^T \Phi(x_i) - y_i)^2 + \lambda \|\beta\|_1.$$

1.1 General feature-based methods

ℓ_2 **extension.** The previous idea can be extended to a general feature-based model:

$$f(x; \beta) = \langle \beta, \Phi(x) \rangle_{\mathcal{H}}, \quad (1.1)$$

where

- \mathcal{H} is the feature space, which can be any Hilbert space;
- $\Phi : \mathcal{X} \mapsto \mathcal{H}$ is the feature map;
- The “coefficients” are $\beta \in \mathcal{H}$.

If taking $\mathcal{H} = \mathbb{R}^m$ and $\Phi(x) = (\phi(x_1), \dots, \phi(x_n))^T \in \mathbb{R}^m$, we recover the classical ones. However, the advantage of the formulation (1.1) is that it includes the case where $m = \infty$. Below is an example:

Random feature models (RFMs). Consider

$$f(x; \beta) = \int \beta(w) \varphi(x; w) d\pi(w) = \langle \beta, \varphi(x; \cdot) \rangle_{L^2(\pi)}, \quad (1.2)$$

where π is a fixed distribution. In this case, the feature map is given by

$$\Phi : \mathcal{X} \mapsto L^2(\pi), \quad \Phi(x) = \varphi(x; \cdot),$$

and the parameter is $\beta \in L^2(\pi)$. The model (1.2) can be viewed as the continuum limit of the following random feature model

$$f(x; \beta) = \frac{1}{m} \sum_{j=1}^m \beta_j \varphi(x; w_j),$$

where w_1, \dots, w_m are independently sampled from π and fixed.

Now, the model (1.1) is well-defined. The objective function of the corresponding ridge regression can be written as

$$\hat{\mathcal{R}}_n(\beta) = \frac{1}{n} \sum_{i=1}^n (\langle \beta, \Phi(x_i) \rangle_{\mathcal{H}} - y_i)^2 + \lambda \|\beta\|_{\mathcal{H}}^2. \quad (1.3)$$

How can we optimize (1.3), which is an infinitely dimensional problem?

1.2 ℓ_1 extension.

Consider the random feature methods:

$$\frac{1}{n} \sum_{i=1}^n \left(\sum_{j=1}^m \beta_j \varphi(x_i; w_j) - y_i \right)^2 + \frac{\lambda}{m} \sum_{j=1}^m |\beta_j|.$$

Assume for any $x \in \mathcal{X}$, $\text{ess sup}_w |\varphi(x; w)| < \infty$. Then, the continuum limit of the above method is given by

$$\min_{\beta \in L^1(\pi)} \frac{1}{n} \sum_{i=1}^n \left(\int \beta(w) \varphi(x_i; w) d\pi(w) - y_i \right)^2 + \lambda \int |\beta(w)| d\pi(w).$$

This method can not be analyzed using the kernel theory.

2 Representer theorem and kernel methods

When it is clear from the context, we will drop the subscripts in $\langle \cdot, \cdot \rangle_{\mathcal{H}}$ and $\| \cdot \|_{\mathcal{H}}$ for simplicity. Let us consider a general problem:

$$\hat{R}_n(\beta) = \frac{1}{2n} \sum_{i=1}^n \ell(f(x_i; \beta), y_i) + \lambda r(\|\beta\|), \quad (2.1)$$

- $f(x; \beta) = \langle \beta, \Phi(x) \rangle$
- ℓ is a general loss function.
- $r : [0, \infty) \mapsto [0, \infty)$ is a strictly increasing function.

Theorem 2.1 (Representer theorem). *Let $\hat{\beta}$ be a minimizer of (2.1). Then, there must exist $a_1, \dots, a_n \in \mathbb{R}$ such that $\hat{\beta} = \sum_{i=1}^n a_i \Phi(x_i)$ and*

$$f(x; \hat{\beta}) = \langle \hat{\beta}, \Phi(x) \rangle = \sum_{i=1}^n a_i k(x_i, x), \quad (2.2)$$

where $k(x, x') := \langle \Phi(x), \Phi(x') \rangle$.

Proof. Let $V_n = \text{span}\{\Phi(x_1), \dots, \Phi(x_n)\} \subset \mathcal{H}$. For any $\beta \in \mathcal{H}$, we can decompose it as follows

$$\beta = \beta_{\parallel} + \beta_{\perp},$$

where $\beta_{\parallel} \in V_n, \beta_{\perp} \in V_n^{\perp}$. Hence, $\|\beta\|^2 = \|\beta_{\parallel}\|^2 + \|\beta_{\perp}\|^2$. Since $r(\cdot)$ is non-decreasing,

$$r(\|\beta\|) \geq r(\|\beta_{\parallel}\|). \quad (2.3)$$

On the other hand, for any x_i ,

$$f(x_i; \beta) = \langle \beta, \Phi(x_i) \rangle = \langle \beta_{\parallel}, \Phi(x_i) \rangle + \langle \beta_{\perp}, \Phi(x_i) \rangle = \langle \beta_{\parallel}, \Phi(x_i) \rangle, \quad (2.4)$$

where the last equality is due to $\beta_{\perp} \in V_n^{\perp}$. Combining (2.3) and (2.4), we have for any $\beta \in \mathcal{H}$,

$$\hat{\mathcal{R}}_n(\beta) \geq \hat{\mathcal{R}}_n(\beta_{\parallel}).$$

Therefore, we can take $\hat{\beta}_{\parallel} = \sum_{i=1}^n a_i \Phi(x_i)$. Then, the function represented can be written as

$$f(x; \beta) = \langle \hat{\beta}_{\parallel}, \Phi(x) \rangle = \sum_{i=1}^n a_i \langle \Phi(x_i), \Phi(x) \rangle = \sum_{i=1}^n a_i k(x_i, x).$$

□

This theorem allows transforming the infinite-dimensional optimization problem (2.1) into a finite dimensional problem. Moreover, we only need to access the kernel $k(\cdot, \cdot)$ without needing to evaluate the feature maps.

The reduced model. Representer theorem implies that we only need to choose

$$\beta = \sum_{j=1}^n a_j \Phi(x_j), \quad f(x; \beta) = \sum_{j=1}^n a_j k(x_j, x).$$

Moreover,

$$\|\beta\|^2 = \left\langle \sum_{j=1}^n a_j \Phi(x_j), \sum_{j=1}^n a_j \Phi(x_j) \right\rangle = \sum_{i,j=1}^n k(x_i, x_j) a_i a_j = a^T K a,$$

where $a = (a_1, \dots, a_n)^T \in \mathbb{R}^n$ and $K = (k(x_i, x_j)) \in \mathbb{R}^{n \times n}$ is the *kernel matrix*.

The kernel ridge regression (KRR) corresponds to the case where $\ell(y, y') = (y - y')^2$ and $r(t) = t^2$, i.e., the problem (1.3). Then, the problem can be reduced to the following n -dimensional problem

$$\hat{\mathcal{R}}_n(a) = \frac{1}{n} \|K a - y\|_2^2 + \lambda a^T K a, \quad (2.5)$$

whose solution is given by

$$a = \left(\frac{1}{n} K + I \right)^{-1} y.$$

In general, kernel methods refer to methods whose hypothesis class is given by

$$\mathcal{F} = \left\{ \sum_{j=1}^n a_j k(x_j, \cdot) : a \in \mathbb{R}^n \right\}.$$

Mathematically, the kernel is defined as follows.

Definition 2.2 (kernel). $k : \mathcal{X} \times \mathcal{X} \mapsto \mathbb{R}$ is said to be a kernel if there exists a feature map $\Phi : \mathcal{X} \mapsto \mathcal{H}$ such that

$$k(x, x') = \langle \Phi(x), \Phi(x') \rangle.$$

Below is a list of popular kernels.

Polynomial kernel: $k(x, x') = (1 + x^T x')^s$ is a kernel for any $s \in \mathbb{N}_+$.

- Linear ($s = 1$). We have $k(x, x') = \langle \Phi(x), \Phi(x') \rangle$ with

$$\Phi(x) = (1, x_1, \dots, x_d).$$

- Quadratic ($s = 2$): The feature map is given by

$$\Phi(x) = \underbrace{(x_d^2, \dots, x_1^2)}_{\text{quadratic}}, \underbrace{(\sqrt{2}x_d x_{d-1}, \dots, \sqrt{2}x_d x_1, \sqrt{2}x_{d-1} x_{d-2}, \dots, \sqrt{2}x_2 x_1)}_{\text{cross terms}}, \underbrace{(\sqrt{2}x_d, \dots, \sqrt{2}x_1)}_{\text{linear terms}}, \underbrace{(1)}_{\text{constant}}).$$

$$\begin{aligned} \langle \Phi(x), \Phi(x') \rangle &= \sum_{i=1}^d (x_i)^2 (x'_i)^2 + 2 \sum_{i \neq j} x_i x_j x'_i x'_j + 2 \sum_i x_i x'_i + 1 \\ &= \left(\sum_{i=1}^d x_i x'_i \right)^2 + 2 \sum_i x_i x'_i + 1 \\ &= (x^T x' + 1)^2 \end{aligned} \quad (2.6)$$

Gaussian kernel: $k(x, x') = e^{-\frac{\|x-x'\|_2^2}{2}}$. Considering $d = 1$, we have

$$\begin{aligned} k(x, x') &= e^{-\frac{x^2}{2} - \frac{x'^2}{2}} e^{xx'} = e^{-\frac{x^2}{2} - \frac{x'^2}{2}} \sum_n \frac{1}{n!} (x)^n (x')^n \\ &= \langle \Phi(x), \Phi(x') \rangle, \end{aligned}$$

where $\Phi(x) = e^{-\frac{x^2}{2}} (1, x, \frac{1}{\sqrt{2}}x^2, \dots, \frac{1}{\sqrt{n!}}x^n, \dots)$.

Laplace kernel:

$$k(x, x') = e^{-\frac{\|x-x'\|_2}{\sigma}}.$$

This kernel is less smooth than the Gaussian kernel. Recently, it has been shown that the Laplace kernel is intimately related to neural network models in the kernel regime.

For a specific problem, choosing appropriate kernels is highly non-trivial. One may need to incorporate the domain knowledge into the kernel design.

3 Reproducing kernel Hilbert spaces

In this section, we ask the question:

What kind of functions can be “efficiently” learned by kernel methods?

By representer theorem, consider

$$\mathcal{F} = \cup_{n=1}^{\infty} \mathcal{F}_n,$$

where

$$\mathcal{F}_n = \left\{ \sum_{j=1}^n a_j k(\cdot, x_j) : x_j \in \mathcal{X}, a_j \in \mathbb{R}, j \in [n] \right\}.$$

This intuition tells us that what kind of functions can be “approximated” by kernel methods. We are interested in functions $f \in \bar{\mathcal{F}}$. However, the problem is how to take the closure and measure the complexity of $f \in \bar{\mathcal{F}}$? Without imposing constraints on the norm of coefficients $\{a_j\}$ in taking the closure, this space can be extremely large. For example, if the corresponding features are polynomials, then \mathcal{F} contains all the continuous functions because of the Stone-Weierstrass theorem. However, $C(\mathcal{X})$ is too large since the Rademacher complexity is $O(1)$. We hope that the Rademacher complexity is on the order of $O(1/\sqrt{n})$.

We need to define an “appropriate” norm for $f \in \mathcal{F}$.

Let us take a step back to the feature-based representation:

$$\beta = \sum_{j=1}^n a_j \Phi(x_j), \quad f(x; \beta) = \sum_{j=1}^n a_j k(x_j, \cdot).$$

In KRR, we penalize $\|\beta\|^2$ of the hypothesis. This means that $\|\beta\|^2$ should be a good norm of the represented function $f(x; \beta)$, i.e.,

$$\|f(\cdot; \beta)\|^2 = \|\beta\|^2 = \left\langle \sum_{i=1}^n a_i \Phi(x_i), \sum_{j=1}^n a_j \Phi(x_j) \right\rangle = \sum_{i,j=1}^n a_i a_j k(x_i, x_j).$$

This intuition can be made rigorous by the following theorem.

Theorem 3.1 (Moore-Aronsjajn theorem). *Let $k : \mathcal{X} \times \mathcal{X} \mapsto \mathbb{R}$ be any kernel. Let $\mathcal{H}^0 = \text{span}(\{k(\cdot, x) : x \in \mathcal{X}\})$ and endow it with the inner product:*

$$\langle f, g \rangle_{\mathcal{H}^0} = \sum_{i=1}^n \sum_{j=1}^m \alpha_i \beta_j k(x_i, x'_j), \quad (3.1)$$

where $f = \sum_{i=1}^n \alpha_i k(\cdot, x_i)$, $g = \sum_{j=1}^m \beta_j k(\cdot, x'_j)$. Then, \mathcal{H}^0 is a valid pre-Hilbert space, i.e, the pointwise closure $\mathcal{H}_k = \overline{\mathcal{H}^0}$ is a Hilbert space.

Proof. We show that (3.1) indeed defines a valid inner product. First,

$$\langle f, g \rangle_{\mathcal{H}^0} = \sum_{i=1}^n \alpha_i g(x_i) = \sum_{j=1}^m \beta_j f(x'_j).$$

It is implied that that the inner product is independent of the specific representation of f and g . The triangular inequality is easy to verify. Next, we show that $\|f\|_{\mathcal{H}^0} = 0$ if and only if $f = 0$. If there exist $x_0 \in \mathcal{X}$ such that $f(x_0) \neq 0$. Assume $f(x) = \sum_{j=1}^m a_j k(x_j, \cdot)$ and consider

$$0 \leq \|\lambda f + f(x_0)k(\cdot, x_0)\|_{\mathcal{H}^0}^2 = \lambda^2 \|f\|_{\mathcal{H}^0}^2 + 2\lambda f^2(x_0) + f^2(x_0)k(x_0, x_0).$$

Taking $\lambda \rightarrow -\infty$, the RHS will be negative and this causes contradictory.

What remains is to show that the convergence of Cauchy sequence. We refer to [Link](#) for a complete proof. \square

Lemma 3.2. *The Hilbert space defined in Theorem 3.1 satisfies the reproducing property:*

$$\langle f, k(\cdot, x) \rangle_{\mathcal{H}_k} = f(x).$$

Proof. For $f \in \mathcal{H}^0$, we can write $f(x) = \sum_{j=1}^m a_j k(\cdot, x_j)$. By definition,

$$\langle f, k(\cdot, x) \rangle_{\mathcal{H}_k} = \sum_{j=1}^m a_j k(x, x_j) = f(x).$$

For any $f \in \mathcal{H}_k$, let $\lim_{n \rightarrow \infty} f_n(x) = f(x)$. Then,

$$\langle f, k(\cdot, x) \rangle_{\mathcal{H}_k} = \lim_{n \rightarrow \infty} \langle f_n, k(\cdot, x) \rangle_{\mathcal{H}_k} = \lim_{n \rightarrow \infty} f_n(x) = f(x).$$

\square

The reproducing property is the most important property of this Hilbert space.

Definition 3.3 (RKHS). Let \mathcal{X} be an arbitrary set and \mathcal{H} a Hilbert space of real-valued functions on \mathcal{X} . We say \mathcal{H} is a reproducing kernel Hilbert space (RKHS) if there is a kernel $k : \mathcal{X} \times \mathcal{X} \mapsto \mathbb{R}$ such that

- $\forall x \in \mathcal{X}, k(\cdot, x) \in \mathcal{H}$.
- *Reproducing property:* $\forall x \in \mathcal{X}, f \in \mathcal{H}, \langle f, k(\cdot, x) \rangle_{\mathcal{H}} = f(x)$

Lemma 3.4. *For a RKHS, the evaluation functional $L_x(f) = f(x)$ is continuous.*

Proof. For any $x \in X$ and $f \in \mathcal{H}$,

$$\sup_{\|f\|_{\mathcal{H}} \leq 1} |L_x(f)| = \sup_{\|f\|_{\mathcal{H}} \leq 1} |\langle f, k(\cdot, x) \rangle_{\mathcal{H}}| \leq \|k(\cdot, x)\|_{\mathcal{H}} < \infty.$$

□

This continuity of the evaluation functional is sometimes used as the equivalent definition of RKHS. An important implication is that the convergence in norm implies the pointwise convergence. If $\lim_{n \rightarrow \infty} \|f_n - f\|_{\mathcal{H}} = 0$, then

$$|f_n(x) - f(x)| \leq \|L_x\| \|f_n - f\|_{\mathcal{H}} \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Lemma 3.5. *For a RKHS, the reproducing kernel k is unique.*

Proof. For any two kernels k_1, k_2 ,

$$\langle f, k_1(\cdot, x) - k_2(\cdot, x) \rangle_{\mathcal{H}} = f(x) - f(x) = 0, \forall x \in X, \forall f \in \mathcal{H}.$$

Taking $f = k_1(\cdot, x) - k_2(\cdot, x)$, we have $\|k_1(\cdot, x) - k_2(\cdot, x)\|_{\mathcal{H}}^2 = 0, \forall x \in X$. Hence, $k_1 = k_2$. □

Theorem 3.6. *For any kernel k , there is a unique RKHS, for which k is the reproducing kernel.*

Proof. First, by Moore-Aronsajn theorem, there exists a RKHS with k being the reproducing kernel. Assume \mathcal{H}_1 and \mathcal{H}_2 be two RKHSs with k being the reproducing kernel. First, by definition, $k(\cdot, x) \in \mathcal{H}_1$ for any $x \in \mathcal{X}$. Hence, $\mathcal{H}^0 \subset \mathcal{H}_1$. Moreover, \mathcal{H}^0 is dense in \mathcal{H}_1 since if there exists $f \in \mathcal{H}$ such that $f \perp \mathcal{H}^0$, we must have

$$\langle f, k(\cdot, x) \rangle_{\mathcal{H}_1} = f(x) = 0 \quad \forall x \in \mathcal{X}.$$

For $f = \sum_{j=1}^m a_j k(\cdot, x_j)$,

$$\begin{aligned} \|f\|_{\mathcal{H}_1}^2 &= \left\langle \sum_{i=1}^n a_i k(\cdot, x_i), \sum_{j=1}^m a_j k(\cdot, x_j) \right\rangle_{\mathcal{H}_1} = \sum_{i,j=1}^n a_i a_j \langle k(\cdot, x_i), k(\cdot, x_j) \rangle_{\mathcal{H}_1} \\ &\stackrel{(i)}{=} \sum_{i,j=1}^n a_i a_j k(x_i, x_j) = \|f\|_{\mathcal{H}^0}^2. \end{aligned}$$

where (i) follows from the reproducing property. Hence, $\|f\|_{\mathcal{H}_1} = \|f\|_{\mathcal{H}^0}$ for $f \in \mathcal{H}^0$. By the same argument, the same results hold for \mathcal{H}_2 . For any $f \in \mathcal{H}_1$, there must exists $(f_n) \subset \mathcal{H}^0$ such that $f(x) = \lim_{n \rightarrow \infty} f_n(x)$. This implies that $f \in \mathcal{H}_2$. Similarly, \mathcal{H}_1 and \mathcal{H}_2 contains the same functions. What remains is to check that the two norms coincide, which results from

$$\|f\|_{\mathcal{H}_1} = \lim_{n \rightarrow \infty} \|f_n\|_{\mathcal{H}_1} = \lim_{n \rightarrow \infty} \|f_n\|_{\mathcal{H}^0} = \lim_{n \rightarrow \infty} \|f_n\|_{\mathcal{H}_2} = \|f\|_{\mathcal{H}_2}.$$

□

Theorem 3.7. *A Hilbert space of functions $\mathcal{H} \subset \mathbb{R}^{\mathcal{X}}$ is a RKHS if and only if the evaluation functional is continuous.*

Proof. If L_x is continuous, by Riesz representation theorem, there exist $K_x \in \mathcal{H}$ such that

$$L_x(f) = \langle K_x, f \rangle_{\mathcal{H}}.$$

Define the kernel:

$$k(x, x') = \langle K_x, K_{x'} \rangle_{\mathcal{H}} = K_{x'}(x) = K_x(x'),$$

for which

$$\langle f, k(\cdot, x) \rangle_{\mathcal{H}} = \langle f, K_x \rangle_{\mathcal{H}} = f(x), \quad \forall f \in \mathcal{H}.$$

This means $k(\cdot, \cdot)$ is a reproducing kernel of \mathcal{H} . □

4 A generalization analysis of kernel ridge regression

We first provide the upper bound of the Rademacher complexity.

Proposition 4.1. *For any kernel k , let \mathcal{H}_k the corresponding RKHS. Let $\mathcal{H}_k^Q = \{f \in \mathcal{H}_k : \|f\|_{\mathcal{H}_k} \leq Q\}$. Then, we have*

$$\widehat{\text{Rad}}_n(\mathcal{H}_k^Q) \leq Q \frac{\sqrt{\sum_{i=1}^n k(x_i, x_i)}}{n}.$$

Proof.

$$\begin{aligned} n\widehat{\text{Rad}}_n(\mathcal{H}_k^Q) &= \mathbb{E}_{\xi} \left[\sup_{\|f\|_{\mathcal{H}_k} \leq Q} \sum_{i=1}^n \xi_i f(x_i) \right] = \mathbb{E}_{\xi} \left[\sup_{\|f\|_{\mathcal{H}_k} \leq Q} \sum_{i=1}^n \xi_i \langle f, k(\cdot, x_i) \rangle_{\mathcal{H}_k} \right] \text{(reproducing property)} \\ &= \mathbb{E}_{\xi} \left[\sup_{\|f\|_{\mathcal{H}_k} \leq Q} \langle f, \sum_{i=1}^n \xi_i k(\cdot, x_i) \rangle_{\mathcal{H}} \right] \leq Q \mathbb{E}_{\xi} \left[\left\| \sum_{i=1}^n \xi_i k(\cdot, x_i) \right\|_{\mathcal{H}_k} \right] \\ &= Q \mathbb{E}_{\xi} \sqrt{\sum_{i,j=1}^n \xi_i \xi_j k(x_i, x_j)} \leq Q \sqrt{\mathbb{E}_{\xi} \left[\sum_{i,j=1}^n \xi_i \xi_j k(x_i, x_j) \right]} \quad \text{(Jensen inequality)} \\ &= Q \sqrt{\sum_{i=1}^n k(x_i, x_i)} \quad (\mathbb{E}[\xi_i \xi_j] = 0, \forall i \neq j). \end{aligned}$$

□

Given data $\{(x_i, f^*(x_i))\}_{i=1}^n$, consider the kernel ridge regression (KRR) estimator

$$\hat{f}_n = \underset{f \in \mathcal{H}_k}{\text{argmin}} \hat{\mathcal{R}}_n(f) + \lambda \|f\|_{\mathcal{H}_k}. \quad (4.1)$$

Theorem 4.2 (A priori estimate). *Assume that $\ell(\cdot, y)$ is L -Lipschitz and bounded by B , and $\sup_{x \in \mathcal{X}} k(x, x) \leq 1$. Then, for any $\delta \in (0, 1)$, with probability $1 - \delta$ over the choice of training set, we have*

$$\mathcal{R}(\hat{f}_n) \lesssim \lambda \|f^*\|_{\mathcal{H}_k} + \frac{L \|f^*\|_{\mathcal{H}_k}}{\sqrt{n}} + B \sqrt{\frac{\log(1/\delta)}{n}}.$$

Proof. (1) Let $Q = \|f^*\|_{\mathcal{H}_k}$. By the definition of \hat{f}_n ,

$$\hat{\mathcal{R}}_n(\hat{f}_n) + \lambda \|\hat{f}_n\|_{\mathcal{H}_k} \leq \hat{\mathcal{R}}_n(f^*) + \lambda \|f^*\|_{\mathcal{H}_k} = \lambda \|f^*\|_{\mathcal{H}_k} = \lambda Q,$$

which yields

$$\|\hat{f}_n\|_{\mathcal{H}_k} \leq Q, \quad \hat{\mathcal{R}}_n(\hat{f}_n) \leq \lambda Q.$$

(2) Let $\mathcal{F}_Q = \{\ell(h(x), h^*(x)) : h \in \mathcal{H}_k^Q\}$. By the contraction lemma, we have

$$\hat{\mathcal{R}}_n(\mathcal{F}_Q) \leq L \hat{\mathcal{R}}_n(\mathcal{H}_k^Q).$$

Using the Rademacher complexity-based generalization bound, we have

$$\begin{aligned} |\hat{\mathcal{R}}_n(\hat{f}_n) - \mathcal{R}(\hat{f}_n)| &\leq \sup_{\|f\|_{\mathcal{H}_k} \leq Q} |\hat{\mathcal{R}}_n(f) - \mathcal{R}(f)| \lesssim \hat{\mathcal{R}}_n(\mathcal{F}_Q) + B \sqrt{\frac{\log(4/\delta)}{n}} \\ &\lesssim L \hat{\mathcal{R}}_n(\mathcal{H}_k^Q) + B \sqrt{\frac{\log(4/\delta)}{n}} \leq \frac{LQ}{\sqrt{n}} + B \sqrt{\frac{\log(1/\delta)}{n}} \quad (\text{use } \sup_{x \in X} k(x, x) \leq 1). \end{aligned}$$

(3) $\mathcal{R}(\hat{f}_n) \leq \hat{\mathcal{R}}_n(\hat{f}_n) + |\hat{\mathcal{R}}_n(\hat{f}_n) - \mathcal{R}(\hat{f}_n)| \leq \lambda Q + (LQ + B \sqrt{\log(4/\delta)})/\sqrt{n}$. □

The preceding estimate is a priori, since it depends on the norm of f^* instead of that of \hat{f}_n . Taking $\lambda = O(1/\sqrt{n})$, we have that $\mathcal{R}(\hat{f}_n) = O(1/\sqrt{n})$, which does not suffer from the curse of dimensionality. This means that the functions in the RKHS can be efficiently learned by the KRR.

- Similar results hold for any regularizations of the form $r(\|f\|_{\mathcal{H}_k})$, where $r : [0, \infty) \rightarrow [0, \infty)$ is strictly increasing.
- Note that Theorem 4.2 holds as long as $\lambda > 0$ and one can even take $\lambda \rightarrow 0^+$, which may seem strange at the first glance. This is due to that there is no label noise. In fact, the optimal λ depends on the level of noise as shown in the following theorem.

Consider the estimator

$$\hat{f}_n = \operatorname{argmin}_{f \in \mathcal{H}_k} \frac{1}{n} \sum_{i=1}^n (T \circ f(x_i) - y_i)^2 + \lambda \|f\|_{\mathcal{H}_k}, \quad (4.2)$$

where $T(t) = \min(\max(t, -1), 1)$. We make the following non-essential technical assumptions.

- $\sup_x |f^*(x)| \leq 1$ and $\sup_x k(x, x) \leq 1$.
- $y_i = f^*(x_i) + \xi_i$. $\{\xi_i\}_i$ are i.i.d. random noises with $|\xi_i| \leq \sigma$. Assume that $\sigma \leq 1$.

Theorem 4.3. *Under the preceding assumptions and taking $\lambda = \frac{\sigma}{\sqrt{n}}$, for any $\delta \in (0, 1)$, we have*

$$\|\hat{f}_n - f^*\|_{L^2(\mathbb{P}_x)}^2 \lesssim \frac{\|f^*\|_{\mathcal{H}_k} + \sqrt{\log(2/\delta)}}{\sqrt{n}}.$$

We refer to [E et al., 2019] for the proof.

Tightness. It is worth noting that preceding bounds are not tight for the square loss: $\ell(y_1, y_2) = (y_1 - y_2)^2$. When applying the contraction lemma, we use the worst-case Lipschitz norm $\text{Lip}(t^2/2) \leq 1$ for $t \in [0, 1]$. However, around the estimator, we should have $\varepsilon(x) = \hat{f}(x) - f^*(x) \ll 1$. Therefore, we should use the “local” Lipschitz norm to bound the Rademacher complexity. This will in turn gives rise to a fast rate. Usually, the fast rate is close to $O(1/n)$ and this approach is called “local Rademacher complexity”. Please refer to [Bartlett et al., 2005] for more details.

References

- [Bartlett et al., 2005] Bartlett, P. L., Bousquet, O., Mendelson, S., et al. (2005). Local Rademacher complexities. *The Annals of Statistics*, 33(4):1497–1537.
- [E et al., 2019] E, W., Ma, C., and Wang, Q. (2019). A priori estimates of the population risk for residual networks. *arXiv preprint arXiv:1903.02154*.